

LYAPUNOV EXPONENTS OF IID LINEAR COCYCLES À LA FURSTENBERG

MARTIM DA COSTA AND JOÃO LOPES DIAS

ABSTRACT. We study the Lyapunov exponents associated to the product of i.i.d. random linear cocycles in $SL(2, \mathbb{R})$. The existence of these quantities and conditions to guarantee strict positivity are established by a celebrated theorem of H. Furstenberg. These results are used to prove the exponential growth of a random Fibonacci sequence and to give an alternative approach to the study of Lévy constants in the context of continued fractions.

1. INTRODUCTION

One of the goals of this work is to provide a self-contained and accessible introduction to the work of Hillel “Harry” Furstenberg in [10, 9], which laid the foundation for the study of Lyapunov exponents of random linear cocycles. This is a topic which has received a lot of recent attention based on its connection to discrete one-dimensional Schrödinger operators. This link with the dynamical systems theory turned out to be very fruitful, leading to many remarkable results, cf. [5, 11, 6].

Lyapunov exponents quantify the exponential norm-growth of a dynamical system. If the Lyapunov exponents are positive, it means that the system displays hyperbolicity and it is often called “chaotic”. This fact reveals much of the asymptotic and statistical behaviour of the orbits. There are very few non-trivial examples for which the positivity of the Lyapunov exponents is known. Furstenberg theorem gives conditions for this property to hold for products of random matrices and linear cocycles.

Our presentation of the proof follows the one by Jairo Bochi [2] which is also inspired in [4]. As such, it is a particular version of the general result, which instead of $SL_{\pm}(2, \mathbb{R})$ considers the larger space $SL_{\pm}(n, \mathbb{R})$. Another goal of this paper is to include the detail and background which is often omitted from the literature, thus providing a gentler presentation of these topics, accessible to a wider audience. This accounts for the rather lengthy preliminary section. Afterwards, we formally state the result and establish useful equivalent conditions. We use these to study the random Fibonacci sequence, and prove that its associated Lyapunov exponent is positive.

Our last goal, in the final section, is to present a novel application of Furstenberg's theorem to a number theoretical problem. We find a direct connection between Lyapunov exponents of random linear cocycles and Lévy constants for the growth of rational approximants in continued fractions. We believe that this relation could be useful to obtain new results. In particular, the recent advances in questions about the regularity of Lyapunov exponents with respect to the distribution of the stochastic iid process underneath, could lead to further insights on the regularity of the Lévy constants.

2. PRELIMINARIES

Let $(\Omega, \mathcal{F}, \rho)$ be a probability space and $f : \Omega \rightarrow \Omega$ a measurable map which preserves ρ , i.e. $f_*\rho = \rho \circ f^{-1} = \rho$ on \mathcal{F} (we also say that ρ is *f-invariant*). Moreover, we denote the set of all 2×2 matrices with determinant ± 1 by $\text{SL}_{\pm}(2, \mathbb{R})$. If $A : \Omega \rightarrow \text{SL}_{\pm}(2, \mathbb{R})$ is measurable, we construct a skew-product map given by

$$\begin{aligned} T : \Omega \times \mathbb{R}^2 &\rightarrow \Omega \times \mathbb{R}^2 \\ (\omega, v) &\mapsto (f(\omega), A(\omega)v). \end{aligned}$$

This is called the *linear cocycle* of A over f , and usually denoted by $T = (f, A)$. The orbit under T of the point $(\omega, v) \in \Omega \times \mathbb{R}^2$ is

$$T^n(\omega, v) = (f^n(\omega), A^{(n)}(\omega)v)$$

where

$$A^{(n)}(\omega) = A(f^{n-1}(\omega))A(f^{n-2}(\omega)) \cdots A(f(\omega))A(\omega).$$

In this paper we are interested in studying one particular linear cocycle. Let (G, \mathcal{X}, μ) be a probability space with $G \subseteq \text{SL}_{\pm}(2, \mathbb{R})$. Define σ to be the shift map

$$\begin{aligned} \sigma : G^{\mathbb{N}} &\rightarrow G^{\mathbb{N}} \\ (\omega_1, \omega_2, \omega_3, \dots) &\mapsto (\omega_2, \omega_3, \dots) \end{aligned}$$

and $A : G^{\mathbb{N}} \rightarrow G$ defined by $(\omega_1, \omega_2, \dots) \mapsto \omega_1$. Both maps are measurable with respect to the infinite-dimensional product space $(G^{\mathbb{N}}, \mathcal{X}^{\mathbb{N}}, \mu^{\mathbb{N}})$, where $\mathcal{X}^{\mathbb{N}}$ denotes the σ -algebra generated by the cylinder sets and $\mu^{\mathbb{N}}$ denotes the product probability measure. The cocycle (σ, A) is called the *product of i.i.d. random matrices*. Its dynamics are given by

$$T^n((\omega_1, \omega_2, \dots), v) = ((\omega_{n+1}, \omega_{n+2}, \dots), \omega_n \cdots \omega_1 v).$$

Throughout this text, M_n will denote the random variable defined by $M_n(\omega) = A(\sigma^{n-1}(\omega)) = \omega_n$.

2.1. Ergodic theory. The map $\tau_v : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by $x \mapsto \tau_v(x) = x + v$ for a fixed $v \in \mathbb{R}^n$ is called a *translation*. The Lebesgue measure λ is known to be the unique measure on \mathbb{R}^n which is τ_v -invariant for all v .

Definition 1. A group (G, \cdot) together with a topology \mathcal{T} is a *topological group* if the maps

$$\begin{aligned} G \times G &\rightarrow G & G &\rightarrow G \\ (x, y) &\mapsto x \cdot y, & x &\mapsto x^{-1} \end{aligned}$$

are continuous.

Definition 2. Let G be a topological group and κ be a measure on $\mathcal{B}(G)$ ¹. The measure κ is said to be:

- *left-translation-invariant* if $\kappa(gA) = \kappa(A)$ for all $A \in \mathcal{B}(G)$ and all $g \in G$;
- *right-translation-invariant* if $\kappa(Ag) = \kappa(A)$ for all $A \in \mathcal{B}(G)$ and all $g \in G$.

Observe that \mathbb{R}^n taken with its usual addition is a topological group. The usual translation of a set $A \subset \mathbb{R}^n$ by a vector v can thus be represented by vA , which is equal to Av by commutativity, so the Lebesgue measure is one example of a measure which is both left- and right-translation-invariant. It is then natural to wonder about the existence of analogous measures in other topological groups.

Theorem 3. *Let G be a compact topological group. There exists a unique probability measure on $\mathcal{B}(G)$ that is both left- and right-translation-invariant. We call it the Haar measure on G .*

Proof. See [17]. □

The following theorem is a fundamental result in ergodic theory. It establishes a connection between the long-term behaviour of a dynamical system and its expected value.

Theorem 4 (Birkhoff's ergodic theorem). *Let (X, \mathcal{X}, κ) be a probability space and $f : X \rightarrow X$ be a measurable map such that κ is f -invariant. If $\phi : X \rightarrow \mathbb{R}$ is κ -integrable, then the limit*

$$\phi_f(x) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \phi(f^k(x))$$

exists for κ -a.e. x and

$$\int_X \phi_f d\kappa = \int_X \phi d\kappa.$$

The sum $\phi(x) + \phi(f(x)) + \cdots + \phi(f^{n-1}(x))$ is called the Birkhoff sum of ϕ .

¹This is the Borel σ -algebra on G .

Proof. See [13]. □

Definition 5. Let (X, \mathcal{X}, κ) be a probability space and $T : X \rightarrow X$ a measurable transformation. The map T is said to be κ -ergodic if for all $A \in \mathcal{X}$

$$T^{-1}(A) = A \implies \kappa(A) = 1 \text{ or } \kappa(A) = 0.$$

Theorem 6 (Kingman's subadditive ergodic theorem). *Let T be a measure preserving transformation on a probability space (X, \mathcal{X}, κ) and $(g_n)_{n \in \mathbb{N}}$ be a sequence of integrable functions which is subadditive, i.e.*

$$g_{n+m}(\omega) \leq g_n(\omega) + g_m(T^n \omega).$$

Then,

$$\lim_{n \rightarrow \infty} \frac{g_n(\omega)}{n} = g(\omega) \in \mathbb{R} \cup \{-\infty\}$$

for κ -a.e. ω , where g is a T -invariant function. If T is ergodic, then g is constant.

Proof. See [18]. □

The proof of the following lemma follows [19] as well as [2].

Proposition 7. *Let (X, κ) be a probability space and $T : X \rightarrow X$ a measurable transformation. Suppose κ is T -invariant and let $f \in L^1(\kappa)$ be a function which satisfies*

$$\lim_{n \rightarrow \infty} \sum_{j=0}^{n-1} f(T^j(x)) = +\infty \quad (2.1)$$

for κ -almost every x . Then

$$\int_X f d\kappa > 0.$$

Proof. Let $(s_n)_{n \in \mathbb{N}}$ be the sequence defined by

$$s_n = \sum_{j=0}^{n-1} f \circ T^j.$$

For $\varepsilon > 0$ we define the two following sets:

$$A_\varepsilon = \{x \in X : \forall n \in \mathbb{N} : s_n(x) \geq \varepsilon\} \text{ and } B_\varepsilon = \bigcup_{k \geq 0} T^{-k}(A_\varepsilon).$$

We begin by proving that

$$\kappa \left(\bigcup_{\varepsilon > 0} B_\varepsilon \right) = 1. \quad (2.2)$$

Suppose that x is such that (2.1) is satisfied and that for every $\varepsilon > 0$ we have $x \notin B_\varepsilon$. In particular $x \notin B_{1/l^2}$ for any $l \geq 1$, i.e. $T^k(x) \notin A_{1/l^2}$

for all $k \geq 0$, or, equivalently, for all $k \geq 0$ there exists $n_l \in \mathbb{N}$ such that $s_{n_l}(T^{k_l}x) < 1/l^2$. Therefore

$$\lim_{l \rightarrow \infty} s_{n_1 + \dots + n_l}(x) < \sum_{l=1}^{\infty} \frac{1}{l^2} = \frac{\pi^2}{6}$$

contradicting (2.1).

Now fix $\varepsilon > 0$ and let $x \in B_\varepsilon$. Then, there exists at least one $k \geq 0$ such that $T^k x \in A_\varepsilon$. Let k_x denote the smallest such k . This entails that

$$\begin{aligned} s_n(T^{k_x}x) &= \sum_{j=0}^{n-1} f(T^j(T^{k_x}x)) \\ &= \sum_{j=k_x}^{n-1} f(T^jx) \\ &\geq \varepsilon \end{aligned}$$

for every $n \geq k_x + 1$. Therefore

$$\sum_{j=0}^{n-1} f(T^jx) \geq \sum_{j=0}^{k_x-1} f(T^jx) + \sum_{j=k_x}^{n-1} \varepsilon \mathbb{1}_{A_\varepsilon}(T^jx). \quad (2.3)$$

Let φ and ψ denote the limit of the Birkhoff averages of f and $\mathbb{1}_{A_\varepsilon}$ respectively. Divide (2.3) by n and then let $n \rightarrow \infty$. We obtain

$$\varphi(x) \geq \varepsilon \psi(x). \quad (2.4)$$

Now note that

$$\begin{aligned} \int \psi(x) d\kappa(x) &= \int \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \mathbb{1}_{A_\varepsilon}(T^jx) d\kappa(x) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \kappa(T^{-j}(A_\varepsilon)) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \kappa(A_\varepsilon) \\ &= \kappa(A_\varepsilon) \end{aligned}$$

and, since $\varphi(x) = 0$ when $x \notin B_\varepsilon$,

$$\begin{aligned} \int \psi(x) d\kappa(x) &= \int_{B_\varepsilon} \psi(x) d\kappa(x) \\ &= \kappa(B_\varepsilon). \end{aligned}$$

By Birkhoff's ergodic Theorem,

$$\int f d\kappa = \int \varphi d\kappa \geq 0$$

so we only need to exclude the case of an equality, which is equivalent to saying $\varphi = 0$ almost everywhere. Assume this is the case. By (2.4) $0 = \varphi(x) \geq \varepsilon \kappa(B_\varepsilon)$ and therefore $\kappa(B_\varepsilon) = 0$ for all ε , which contradicts (2.2). The desired result follows from this contradiction. \square

2.2. Lyapunov exponents. *Lyapunov exponents* are quantities associated to a linear cocycle. For a linear cocycle (f, A) , its (upper) Lyapunov exponent γ is defined as

$$\gamma = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|A^{(n)}\|.$$

If the cocycle in question is the product of 2×2 random matrices, which we intend to study, we obtain

$$\gamma = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|M_n \cdots M_1\|. \quad (2.5)$$

Suppose such a quantity exists and consider another, arbitrary norm $\|\cdot\|_*$ on $\mathbb{R}^{2 \times 2}$. From the finite dimension of $\mathbb{R}^{2 \times 2}$ it follows that any two norms are equivalent. Therefore there exists a pair of real numbers $0 < C_1 < C_2$ such that the following inequality is satisfied

$$C_1 \|M_n \cdots M_1\| \leq \|M_n \cdots M_1\|_* \leq C_2 \|M_n \cdots M_1\|.$$

The logarithm function preserves the inequalities. We can then divide by n and take the limit to obtain

$$\gamma = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|M_n \cdots M_1\|_*.$$

This establishes that γ does not depend on the chosen norm, supposing its existence. We now turn to the question of whether or not γ is well-defined.

For an arbitrary function g define g^+ to be the map $x \mapsto \sup(f(x), 0)$. If $\log^+ \|M_1\|$ is integrable and $n, p \geq 1$ then

$$\begin{aligned} \log \|A^{(n+p)}(\omega)\| &\leq \log \|M_{n+p}(\omega) \cdots M_{n+1}(\omega)\| + \log \|M_n(\omega) \cdots M_1(\omega)\| \\ &= \log \|A^{(p)}(\sigma^n(\omega))\| + \log \|A^{(n)}(\omega)\| \end{aligned}$$

by the submultiplicative property of matrix norms. Consequently the sequence $(\log \|A^{(n)}\|)_{n \in \mathbb{N}}$ is subadditive and integrable. By Theorem 6 we have

$$\frac{1}{n} \log \|A^{(n)}(\omega)\| \rightarrow \gamma(\omega) \in \mathbb{R} \cup \{-\infty\}$$

for μ -a.e. $\omega \in \Omega^{\mathbb{N}}$. Since σ is $\mu^{\mathbb{N}}$ -ergodic, γ is almost surely constant. This establishes the conditions for the existence of the Lyapunov exponent γ .

Unless stated otherwise, we fix $\|M\|$ to be the spectral norm of a matrix M , i.e. the square root of the maximum eigenvalue of $M^T M$ and $\|v\|$ the usual euclidean norm for a vector $v \in \mathbb{R}^2$.

2.2.1. Examples.

Example 1. Consider a probability space $(\Omega, \mathcal{F}, \mu)$ such that $\text{supp}(\mu) = O(2)^2$. We have

$$\begin{aligned} \gamma &= \int_{O(2)^{\mathbb{N}}} \lim_{n \rightarrow \infty} \frac{1}{n} \log \|M_n(\omega) \cdots M_1(\omega)\| d\mu^{\mathbb{N}}(\omega) \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{n} \log(1) \\ &= 0 \end{aligned}$$

using the fact that the product $M_n(\omega) \cdots M_1(\omega)$ is an element of $O(2)$ by closure of the group operation, so $\|M_n(\omega) \cdots M_1(\omega)\| = 1$.

Example 2. Suppose

$$\text{supp}(\mu) = \left\{ \begin{bmatrix} t & 0 \\ 0 & 1/t \end{bmatrix} : t \geq 1 \right\}.$$

Observe that

$$M_n \cdots M_1 = \begin{bmatrix} t_n & 0 \\ 0 & 1/t_n \end{bmatrix} \cdots \begin{bmatrix} t_1 & 0 \\ 0 & 1/t_1 \end{bmatrix} = \begin{bmatrix} t_n \cdots t_1 & 0 \\ 0 & (t_n \cdots t_1)^{-1} \end{bmatrix}.$$

So $\|M_n \cdots M_1\| = \max\{t_n \cdots t_1, (t_n \cdots t_1)^{-1}\} = t_n \cdots t_1$. Hence,

$$\log(\|M_n \cdots M_1\|) = \log(t_n \cdots t_1).$$

By the usual law of large numbers

$$\gamma = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \log(t_i) = \mathbb{E} [\log \|M_1\|] > 0.$$

2.3. The one-dimensional projective space. We define the real projective space of dimension one by first defining an equivalence relation \sim on $\mathbb{R}^2 \setminus \{0\}$, stipulating that $x \sim y$ iff there exists $\alpha \in \mathbb{R}$ such that $x = \alpha y$. The real projective space is defined as the quotient

$$\mathbb{RP}^1 = \mathbb{R}^2 \setminus \{0\} / \sim,$$

i.e. the set of all equivalence classes. The equivalence class, or *direction*, of $x \in \mathbb{R}^2 \setminus \{0\}$ will be denoted by \bar{x} and may be thought of as a straight line passing through the origin or as the set of all linear combinations of x denoted by $\text{span}\{x\}$. Such lines are entirely characterized by the angle they form with the horizontal axis. Therefore there is an intuitive identification between \mathbb{RP}^1 and the interval $[0, \pi)$ and the two sets may be regarded as interchangeable when convenient.

A matrix A in $\text{GL}(2, \mathbb{R})$ induces a transformation on \mathbb{RP}^1 in a straightforward manner: we start with an element $\bar{x} \in \mathbb{RP}^1$ and consider $x \in \bar{x}$. We then perform the standard matrix multiplication Ax and take the equivalence class \overline{Ax} . This procedure results in a well-defined function, given that it does not depend on the choice of x . In other words, for

²For a brief discussion of $O(2)$ see section 2.4.

a given matrix $A \in \text{GL}(2, \mathbb{R})$, the induced map $\bar{A} : \mathbb{RP}^1 \rightarrow \mathbb{RP}^1$, is defined by $\bar{x} \mapsto \overline{Ax}$. We selectively adhere to this notational distinction between a matrix and the map it induces in the projective space, and likewise for a vector and its corresponding direction. More often than not, we simply use the same symbol for both, unless we deem it confusing.

The following lemma will be used later on.

Lemma 8. *If $A \in \mathbb{R}^{2 \times 2}$ has rank one and ν is a probability measure on \mathbb{RP}^1 , then $\bar{A}_*\nu$ is a Dirac measure.*

Proof. Since $\text{rank}(A) = 1$, then there exists $x \in \mathbb{R}^2$ such that $\text{range } A = \text{span}(x) = \bar{x}$. Let $B \subseteq \mathbb{RP}^1$ be a measurable set. Then

$$\bar{A}_*\nu(B) = \nu(\bar{A}^{-1}(B)) = \begin{cases} \nu(\mathbb{RP}^1) & \text{if } \bar{x} \in B \\ \nu(\emptyset) & \text{otherwise.} \end{cases}$$

Therefore $\bar{A}_*\nu = \delta_{\bar{x}}$. □

Finally, we note that \mathbb{RP}^1 is a compact and separable topological space.

2.4. Orthogonal matrices. A matrix $A \in \mathbb{R}^{n \times n}$ is said to be *orthogonal* if $AA^\top = I$. We denote the set of all orthogonal $n \times n$ matrices by $O(n)$, which is easily checked to be a subgroup of $\text{GL}(n, \mathbb{R})$.

Suppose $A \in O(n)$. Since $AA^\top = I$ implies that

$$\det(AA^\top) = \det(A) \det(A^\top) = (\det A)^2 = 1$$

then $|\det A| = 1$, or, equivalently, $A \in \text{SL}_\pm(n, \mathbb{R})$.

We now particularize our discussion to 2×2 matrices. Consider

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

an arbitrary element of $O(2)$. Since A^\top is an invertible matrix, its corresponding linear transformation $R_{A^\top} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is an isomorphism, hence

$$\text{span}\{(a, b), (c, d)\} = \text{range } R_{A^\top} = \mathbb{R}^2,$$

from which it follows that the rows of the matrix A form a basis of \mathbb{R}^2 . Additionally, it is an orthonormal basis, because

$$AA^\top = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} a & c \\ b & d \end{bmatrix} = \begin{bmatrix} \|(a, b)\|^2 & (a, b) \cdot (c, d) \\ (a, b) \cdot (c, d) & \|(c, d)\|^2 \end{bmatrix} = I.$$

The fact that $\|(a, b)\| = \|(c, d)\| = 1$ implies the existence of $\theta \in [0, 2\pi)$ such that $(a, b) = (\cos \theta, \sin \theta)$ and, since $(a, b) \perp (c, d)$, the vector (c, d) is either $(-\sin \theta, \cos \theta)$ or $(\sin \theta, -\cos \theta)$. We have proven that

$$O(2) = \left\{ \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} : \theta \in [0, 2\pi) \right\} \cup \left\{ \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{bmatrix} : \theta \in [0, 2\pi) \right\}.$$

Therefore, an arbitrary matrix $A \in O(2)$ has norm equal to $\|A\| = 1$.

3. THE STATEMENT OF THE THEOREM

We now deal exclusively with the product of random matrices, the linear cocycle (σ, A) described earlier and its associated Lyapunov exponent. We assume that the measure μ is such that the associated Lyapunov exponent γ exists, i.e. we assume that $\log^+ \|M\|$ is integrable.

Theorem 9 (Furstenberg). *Let G_μ be the smallest closed subgroup which contains the support of μ . Assume that:*

- i) G_μ is not compact.*
- ii) For every finite, non-empty $L \subseteq \mathbb{RP}^1$, there exists $M \in G_\mu$ such that $\bar{M}(L) \neq L$.*

Then $\gamma > 0$.

The next two propositions establish equivalent conditions for the theorem, which are, in practice, simpler to check.

Proposition 10. *G_μ is compact iff there exists $C \in \text{GL}(2, \mathbb{R})$ such that $CMC^{-1} \in O(2)$ for every $M \in G_\mu$.*

Proof. We begin by proving that

$$\exists C \in \text{GL}(2, \mathbb{R}), \forall M \in G_\mu : CMC^{-1} \in O(2) \implies G_\mu \text{ is compact.}$$

Assume that the premise of the implication above holds and let A be an element of G_μ . Then $CAC^{-1} \in O(2)$ and there exists $R \in O(2)$ such that $A = C^{-1}RC$. Applying the norm to both sides of this equality yields

$$\|A\| = \|C^{-1}RC\| \leq \|C^{-1}\| \|R\| \|C\| = \|C^{-1}\| \|C\|.$$

Since the matrix C is fixed and our choice for A was arbitrary, this argument holds for all elements of G_μ . Consequently, G_μ is bounded and therefore it is compact.

We still have to prove the converse implication. To this end, suppose G_μ is compact. By Theorem 3, a probability measure h , known as Haar measure, exists on G_μ which is both left- and right-translation-invariant. Define the quadratic form $Q_0 : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $Q_0(v) = v^\top I v$ and $Q : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$Q(v) = \int_{G_\mu} Q_0(gv) dh(g) = \int_{G_\mu} \|gv\|^2 dh(g) \geq 0$$

which is a positive quadratic form. There exists a positive semidefinite matrix B such that $Q(v) = v^\top B v$ for every v . So $B = C^\top C$ for some matrix C and

$$Q_0(Cv) = v^\top C^\top C v = Q(v). \quad (3.1)$$

Note that C is invertible. Now let $T_{g_0} : G_\mu \rightarrow G_\mu$ be a translation map defined by $g \mapsto gg_0$, then

$$\begin{aligned} Q(v) &= \int_{G_\mu} Q_0(gv) dh(g) \\ &= \int_{T_{g_0}^{-1}(G_\mu)} Q_0(gv) dh(g) \\ &= \int_{G_\mu} Q_0(gg_0v) dh(g) \\ &= Q(g_0v). \end{aligned}$$

This, together with (3.1), yields

$$Q_0(Cg_0C^{-1}w) = Q_0(w)$$

which means $Cg_0C^{-1} \in O(2)$ as desired. \square

Before we prove the other equivalence we referred to, we need a technical lemma.

Lemma 11. *If $M \in \mathrm{SL}_\pm(2, \mathbb{R})$ fixes three directions then $M = \pm I$.*

Proof. Let $M \in \mathrm{SL}_\pm(2, \mathbb{R})$. Suppose there exist distinct $\bar{x}_1, \bar{x}_2, \bar{x}_3 \in \mathbb{RP}^1$ such that $\bar{M}(\bar{x}_i) = \bar{x}_i$ for $i = 1, 2, 3$. Equivalently, there exists $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ such that $Mx_i = \lambda_i x_i$. The matrix M has at most two linearly independent eigenvectors, thus, without loss of generality, suppose

$$x_3 = \alpha x_1 + \beta x_2$$

for some $\alpha, \beta \in \mathbb{R} \setminus \{0\}$. Then

$$Mx_3 = \alpha Mx_1 + \beta Mx_2 = \alpha \lambda_1 x_1 + \beta \lambda_2 x_2$$

and

$$\lambda_3 x_3 = \alpha \lambda_1 x_1 + \beta \lambda_2 x_2.$$

By linear independence $\lambda_1 = \lambda_2 = \lambda_3$. Additionally, $|\det M| = \lambda_1^2 = 1$ implies that $\lambda_1 \in \{-1, 1\}$ and therefore $M = \pm I$ as desired. \square

We will also need the following proposition, which we state without proof.

Proposition 12. *Let $\varphi: G \rightarrow G'$ be a group homomorphism. The quotient group $G/\ker \varphi$ is isomorphic to $\mathrm{Im} \varphi$.*

Proof. See [1]. \square

Proposition 13. *Assume G_μ is not compact. Condition ii) in Theorem 9 is true iff for every set $L \subseteq \mathbb{RP}^1$ with $\#L \in \{1, 2\}$ there exists $M \in G_\mu$ such that $\bar{M}(L) \neq L$.*

Proof. The \implies direction is trivial. We prove the converse. Suppose $L \subseteq \mathbb{R}\mathbb{P}^1$ is finite, i.e. $L = \{\bar{x}_1, \dots, \bar{x}_n\}$ and $\#L = n$. By hypothesis

$$\bar{M}(L) = \{\bar{M}(\bar{x}_1), \bar{M}(\bar{x}_2), \dots, \bar{M}(\bar{x}_n)\} = L.$$

Since $\#M(L) = \#L$, each matrix $M \in G_\mu$ induces a permutation φ_M of L . This allows us to define a group homomorphism $\varphi : G_\mu \rightarrow \text{Perm}(L)$ where $\text{Perm}(L)$ denotes the group of all permutations of L . The group $\text{Perm}(L)$ is finite and G_μ must be infinite since we are assuming it is non-compact. By Proposition 12, $G_\mu/\text{Ker } \varphi$ is isomorphic to $\text{Im } \varphi \subseteq \text{Perm}(L)$ and therefore $G_\mu/\text{Ker } \varphi$ is also finite. Since

$$G_\mu = \bigcup_{H \in \frac{G_\mu}{\text{Ker } \varphi}} H \text{ker } \varphi$$

is a finite union, each class in $G_\mu/\text{Ker } \varphi$ must be infinite. Consequently, $\text{Ker } \varphi = \{M \in G_\mu : \varphi(M) = I\} = \{M \in G_\mu : M(x_i) = x_i \text{ for } i = 1, \dots, n\}$ is an infinite set. If $n \geq 3$ then, by Lemma 11, $\text{Ker } \varphi$ is finite. This is a contradiction and therefore $n \in \{1, 2\}$. \square

3.1. An application to the random Fibonacci sequence.

Example 3. Consider the random Fibonacci sequence

$$F_n = \begin{cases} F_{n-1} + F_{n-2}, & \text{with probability } p \\ F_{n-1} - F_{n-2}, & \text{with probability } 1 - p \end{cases}$$

for $n \geq 2$ and $F_0 = 0, F_1 = 1$. The classical Fibonacci sequence occurs when $p = 1$, and in this case F_n grows exponentially. We would like to see how F_n evolves when $0 < p < 1$. Notice that

$$\begin{bmatrix} F_{n+2} \\ F_{n+1} \end{bmatrix} = \begin{bmatrix} 1 & \pm 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix}.$$

Define

$$A_+ = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \text{ and } A_- = \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}.$$

We consider the probability space $(\Omega, \mathcal{F}, \mu)$ where $\Omega = \{A_+, A_-\}$, $\mathcal{F} = \mathcal{P}(\Omega)$ and $\mu = p\delta_{A_+} + (1-p)\delta_{A_-}$ with $0 < p < 1$.

In this case, the product of random matrices cocycle will be over the product space $(\Omega^{\mathbb{N}}, \mathcal{F}^{\mathbb{N}}, \mu^{\mathbb{N}})$.

Let G_μ be the smallest closed group which contains $\text{supp}(\mu) = \Omega$. We denote the classical Fibonacci sequence by $(C_n)_{n \in \mathbb{N}}$. Then

$$A_+^n = \begin{bmatrix} C_n & C_{n-1} \\ C_{n-1} & C_{n-2} \end{bmatrix}$$

for $n \geq 2$. Given that C_n grows exponentially, we have $\|A_+^n\| \rightarrow \infty$ as $n \rightarrow \infty$ and this proves that G_μ is not compact. We next check if the second condition of Theorem 9 is satisfied.

If L is made up of only one direction, then the fact that A_- has no real eigenvalues shows that the condition is satisfied.

Suppose $L = \{\bar{x}_1, \bar{x}_2\}$ with $\bar{x}_1 \neq \bar{x}_2$ and $\bar{M}(L) = L$ for every $M \in G_\mu$. The matrix A_- cannot fix both directions since this would again imply it has real eigenvalues. The remaining case is

$$\bar{A}_- \bar{x}_1 = \bar{x}_2 \text{ and } \bar{A}_- \bar{x}_2 = \bar{x}_1$$

which implies $\bar{A}_-^2 \bar{x}_i = \bar{x}_i$ for $i \in \{1, 2\}$. Since \bar{A}_-^2 has complex eigenvalues, no such set L exists. By Proposition 13 and Theorem 9, the associated Lyapunov exponent γ is positive.

For a computer assisted computation of the Lyapunov exponent in this context see [21].

4. PROOF OF THE THEOREM

In this section we fix μ to be a measure satisfying the assumptions of Theorem 9. As mentioned in the introduction, the proof follows the one in the notes by Jairo Bochi [2].

4.1. Properties of measures. Let (X, \mathcal{X}, κ) be a measure space. The measure κ is said to be *atomic* if there exists $x \in X$ such that $\kappa(\{x\}) \neq 0$. A well known example of an atomic measure is the Dirac measure.

If X is a topological space, we denote the space of all the probability measures on $(X, \mathcal{B}(X))$ by the symbol $\mathcal{M}(X)$ endowed with the weak* topology. A detailed study of this topology is beyond the scope of this text, but we will make use of the fact that $\mathcal{M}(X)$ is a compact space if X is compact. This is the case when $X = \mathbb{RP}^1$. Furthermore, weak* convergence is equivalent to the usual weak convergence of measures, that is, (ρ_n) converges to ρ iff

$$\lim_{n \rightarrow \infty} \int f(x) d\rho_n(x) = \int f(x) d\rho(x)$$

for every bounded and continuous function f on X . In this case we write $\rho_n \Rightarrow \rho$. Further details and proofs of these results can be found in [20].

Lemma 14. *If $\nu \in \mathcal{M}(\mathbb{RP}^1)$ is non-atomic and $(A_n \neq 0)_{n \in \mathbb{N}}$ is a sequence of matrices converging to $A \neq 0$, then $A_{n*} \nu \Rightarrow A_* \nu$.*

Proof. Let $f: \mathbb{R}\mathbb{P}^1 \rightarrow \mathbb{R}$ be a continuous function, then

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}\mathbb{P}^1} f(x) dA_{n*}\nu(x) &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}\mathbb{P}^1} f \circ A_n(x) d\nu(x) \\ &= \int_{\mathbb{R}\mathbb{P}^1} \lim_{n \rightarrow \infty} f \circ A_n(x) d\nu(x) \\ &= \int_{\mathbb{R}\mathbb{P}^1} f \circ A(x) d\nu(x) \\ &= \int_{\mathbb{R}\mathbb{P}^1} f(x) dA_*\nu(x), \end{aligned}$$

where we have used the dominated convergence theorem and the continuity of f . \square

Whenever convenient we may omit the domain of integration. In this case, the reader should assume that the domain is the whole sample space corresponding to the respective measure.

Lemma 15. *If $\nu \in \mathcal{M}(\mathbb{R}\mathbb{P}^1)$ is non-atomic, then the set of matrices which preserve ν , i.e.*

$$H_\nu = \{M \in \mathrm{SL}_\pm(2, \mathbb{R}) : M_*\nu = \nu\},$$

is a compact subgroup of $\mathrm{SL}_\pm(2, \mathbb{R})$.

Proof. It is a simple exercise in algebra to see that H_ν is a group. We prove compactness. H_ν is compact iff it is closed and bounded. Let $(M_n \in H_\nu)_{n \in \mathbb{N}}$ be a sequence converging to $M \in \mathbb{R}^{2 \times 2}$. For each n we have $\det M_n = \pm 1$ which implies $M_n \neq 0$. As for the matrix M , note that

$$\det(M) = \det\left(\lim_{n \rightarrow \infty} M_n\right) = \lim_{n \rightarrow \infty} \det(M_n) = \pm 1,$$

and therefore $M \neq 0$. We can apply Lemma 14 to obtain $M_{n*}\nu \Rightarrow M_*\nu$ i.e. $\nu \Rightarrow M_*\nu$ and $\nu = M_*\nu$ as desired. We have proven that H_ν is a closed set.

Suppose, in order to arrive at a contradiction, that H_ν is not bounded, so there exists a sequence $(M_n \in H_\nu)_{n \in \mathbb{N}}$ which diverges. Consider the new sequence (X_n) given by $X_n = M_n \|M_n\|^{-1}$. Since (X_n) is a sequence in a compact subspace of $\mathbb{R}^{2 \times 2}$ it has a convergent subsequence (X_{n_k}) with limit C . Again, by Lemma 14, $X_{n_k*}\nu \Rightarrow C_*\nu$ such that $\nu = C_*\nu$. Now note that,

$$\det C = \det\left(\lim_{k \rightarrow \infty} \frac{M_{n_k}}{\|M_{n_k}\|}\right) = \lim_{k \rightarrow \infty} \frac{\pm 1}{\|M_{n_k}\|^2} = 0.$$

By the fundamental theorem of linear maps, $\mathrm{rank}(C) = 1$. Lemma 8 would then imply that ν is an atomic measure, contradicting our assumption. \square

4.2. Stationary measures.

Definition 16. Let $\nu \in \mathcal{M}(\mathbb{RP}^1)$. We define $\mu * \nu$ to be the measure on \mathbb{RP}^1 which satisfies

$$\int f(x) d(\mu * \nu)(x) = \iint f(\bar{M}x) d\mu(M) d\nu(x)$$

for any bounded Borel function $f : \mathbb{RP}^1 \rightarrow \mathbb{R}$. The measure ν is said to be μ -stationary if $\mu * \nu = \nu$.

We define the evaluation map by

$$\begin{aligned} \text{ev} : \text{SL}_{\pm}(2, \mathbb{R}) \times \mathbb{RP}^1 &\rightarrow \mathbb{RP}^1 \\ (M, \bar{v}) &\mapsto \bar{M}\bar{v}. \end{aligned}$$

Let $\nu \in \mathcal{M}(\mathbb{RP}^1)$. Notice that if $B \subseteq \mathbb{RP}^1$ is a measurable set, then

$$\begin{aligned} \mu * \nu(B) &= \iint \mathbb{1}_B(\text{ev}(A, \bar{x})) d\mu(A) d\nu(\bar{x}) \\ &= \int \mathbb{1}_{\text{ev}^{-1}(B)}(A, \bar{x}) d(\mu \times \nu)(A, \bar{x}) \\ &= \mu \times \nu(\{(A, \bar{x}) : \bar{A}\bar{x} \in B\}) \\ &= (\mu \times \nu)(\text{ev}^{-1}(B)) \\ &= \text{ev}_*(\mu \times \nu)(B). \end{aligned}$$

Lemma 17. *Every μ -stationary $\nu \in \mathcal{M}(\mathbb{RP}^1)$ is non-atomic.*

Proof. Suppose, so as to obtain a contradiction, that ν is atomic. Then, the quantity

$$\beta = \max_{x \in \mathbb{RP}^1} \nu(\{x\})$$

is positive. Let $L = \{x \in \mathbb{RP}^1 : \nu(\{x\}) = \beta\}$. If L has infinite cardinality, then we may consider a countable subset $L_1 = \{x_1, x_2, \dots\}$, but this contradicts the assumption that ν is a probability measure since

$$\begin{aligned} \nu(L_1) &= \nu(\{x_1, x_2, \dots\}) \\ &= \sum_{i=1}^{\infty} \beta \\ &= \infty. \end{aligned}$$

Consequently, L must be finite. Now let $x_0 \in L$ and note that

$$\beta = \nu(\{x_0\}) = \iint \mathbb{1}_{\{M^{-1}x_0\}}(x) d\nu(x) d\mu(M) = \int \nu(\{M^{-1}x_0\}) d\mu(M) \leq \beta$$

By definition, the inequality $\beta \geq \nu(\{M^{-1}x_0\})$ is true for every M , so $\nu(\{M^{-1}x_0\}) = \beta$ and thus $M^{-1}x_0 \in L$ for μ -a.e. M , i.e. $M(L) = L$ for μ -a.e. M . This means that the set

$$F_L = \{M \in \mathrm{SL}_{\pm}(2, \mathbb{R}) : M(L) = L\}$$

has full measure, i.e. $\mu(F_L) = 1$. Furthermore, F_L is closed, so $\mathrm{supp}(\mu) \subseteq F_L$, which implies that $G_{\mu} \subseteq F_L$. This contradicts assumption ii) of Theorem 9. \square

Remark 18. It can be proven that μ -stationary measures always exist (see Lemma 3.5 of [4]). By Lemma 17, any such measure on \mathbb{RP}^1 is non-atomic.

4.3. Convergence of μ -stationary measures. Let $S_n = M_1 \cdots M_n$.

Lemma 19. *Let $\nu \in \mathcal{M}(\mathbb{RP}^1)$ be μ -stationary. For $\mu^{\mathbb{N}}$ -a.e. $\omega \in \Omega$, there exists $\nu_{\omega} \in \mathcal{M}(\mathbb{RP}^1)$ such that*

$$S_n(\omega)_* \nu \Rightarrow \nu_{\omega}.$$

Proof. Let $f \in C(\mathbb{RP}^1)$. Define

$$\begin{aligned} F_f : \mathrm{SL}_{\pm}(2, \mathbb{R}) &\rightarrow \mathbb{R} \\ M &\mapsto \int f(Mx) d\nu(x). \end{aligned}$$

Let \mathcal{F}_n be the σ -algebra of $\mathrm{SL}_{\pm}(2, \mathbb{R})^{\mathbb{N}}$ formed by the cylinders of length n . Then $S_n(\cdot)$ is \mathcal{F}_n -measurable. Let $C \in \mathcal{F}_n$, then

$$\int_C \int_{\mathrm{SL}_{\pm}(2, \mathbb{R})} F_f(S_n(\omega)M) d\mu(M) d\mu^{\mathbb{N}}(\omega) = \int_C F_f(S_{n+1}(\omega)) d\mu^{\mathbb{N}}(\omega).$$

By definition of conditional expectation, we obtain

$$\begin{aligned} \mathbb{E}[F_f(S_{n+1}) \mid \mathcal{F}_n] &= \int_{\mathrm{SL}_{\pm}(2, \mathbb{R})} F_f(S_n M) d\mu(M) \\ &= \iint f(S_n Mx) d\nu(x) d\mu(M) \\ &= \int f(S_n y) d\nu(y) \\ &= F_f(S_n). \end{aligned}$$

Therefore the stochastic process $\{F_f(S_n)\}_{n \in \mathbb{N}}$ is a bounded martingale and as such it converges almost surely, i.e. the limit

$$\Gamma f(\omega) = \lim_{n \rightarrow \infty} F_f(S_n(\omega))$$

exists for a.e. $\omega \in \Omega^{\mathbb{N}}$. We now use this fact to prove $S_n(\omega)_* \nu \Rightarrow \nu_{\omega}$ almost surely for some $\nu_{\omega} \in \mathcal{M}(\mathbb{RP}^1)$.

By the compactness of \mathbb{RP}^1 , the space $C(\mathbb{RP}^1)$ is separable. Let $\{f_k\}_{k \in \mathbb{N}}$ be a dense subset of $C(\mathbb{RP}^1)$. The limit $\Gamma f_k(\omega)$ exists in a set \mathcal{L}_k of full measure for each $k \in \mathbb{N}$. Let

$$\mathcal{L} = \bigcap_{k \in \mathbb{N}} \mathcal{L}_k$$

then

$$\mu^{\mathbb{N}}(\mathcal{L}^{\mathbb{C}}) = \mu^{\mathbb{N}}\left(\bigcup_{n \in \mathbb{N}} \mathcal{L}_k^{\mathbb{C}}\right) \leq \sum_{n=1}^{\infty} \mu^{\mathbb{N}}(\mathcal{L}_k^{\mathbb{C}}) = 0$$

and we conclude that $\mu^{\mathbb{N}}(\mathcal{L}) = 1$. Now consider $\omega \in \mathcal{L}$ and let ν_{ω} be a weak* limit point of the sequence of measures $S_n(\omega)_* \nu$. Then

$$\begin{aligned} \int f_k d\nu_{\omega} &= \lim_{n \rightarrow \infty} \int f_k dS_n(\omega)_* \nu \\ &= \lim_{n \rightarrow \infty} \int f_k \circ S_n(\omega) d\nu \\ &= \lim_{n \rightarrow \infty} F_{f_k}(S_n(\omega)) d\nu \\ &= \Gamma f_k(\omega). \end{aligned}$$

Since the limit is the same for all subsequences then $S_n(\omega)_* \nu \Rightarrow \nu_{\omega}$. \square

Lemma 20. *The measures ν and ν_{ω} from Lemma 19 satisfy*

$$S_n(\omega)_* M_* \nu \Rightarrow \nu_{\omega} \quad \text{as } n \rightarrow \infty$$

for μ -a.e. M .

Proof. Let $\ell = \{f_1, f_2, \dots\}$ be a countable dense subset of $C(\mathbb{R}P^1)$ and fix $k \in \mathbb{N}$. We will prove that the following quantity is finite:

$$I = \int \mathbb{E}^{\mu^{\mathbb{N}}} \left[\sum_{n=1}^{\infty} \left(\int f_k(S_n(\omega) M x) d\nu(x) - \int f_k(S_n(\omega) x) d\nu(x) \right)^2 \right] d\mu(M).$$

Note that

$$\begin{aligned} I &= \int \sum_{n=1}^{\infty} \mathbb{E}^{\mu^{\mathbb{N}}} \left[\left(\int f_k(S_n(\omega) M x) d\nu(x) - \int f_k(S_n(\omega) x) d\nu(x) \right)^2 \right] d\mu(M) \\ &= \sum_{n=1}^{\infty} \int \mathbb{E}^{\mu^{\mathbb{N}}} \left[\left(\int f_k(S_n(\omega) M x) d\nu(x) - \int f_k(S_n(\omega) x) d\nu(x) \right)^2 \right] d\mu(M). \end{aligned}$$

Define

$$\begin{aligned} I_n &= \int \mathbb{E}^{\mu^{\mathbb{N}}} \left[\left(\int f_k(S_n(\omega) M x) d\nu(x) - \int f_k(S_n(\omega) x) d\nu(x) \right)^2 \right] d\mu(M) \\ &= \int \mathbb{E}^{\mu^{\mathbb{N}}} [(F_{f_k}(S_n(\omega) M) - F_{f_k}(S_n(\omega)))^2] d\mu(M) \\ &= \int \mathbb{E}^{\mu^{\mathbb{N}}} [(F_{f_k}(S_n(\omega) M))^2 + (F_{f_k}(S_n(\omega)))^2 - 2F_{f_k}(S_n(\omega) M)F_{f_k}(S_n(\omega))] d\mu(M) \\ &= \mathbb{E}^{\mu^{\mathbb{N}}} \left[\int (F_{f_k}(S_n(\omega) M))^2 + (F_{f_k}(S_n(\omega)))^2 - 2F_{f_k}(S_n(\omega) M)F_{f_k}(S_n(\omega)) d\mu(M) \right] \\ &= \mathbb{E}^{\mu^{\mathbb{N}}} [(F_{f_k}(S_{n+1}(\omega)))^2 + (F_{f_k}(S_n(\omega)))^2 - 2F_{f_k}(S_{n+1}(\omega))F_{f_k}(S_n(\omega))], \end{aligned}$$

where the last equality comes from the fact that

$$\iint F_{f_k}(S_n(\omega)M)^2 d\mu(M)d\mu^{\mathbb{N}}(\omega) = \int F_{f_k}(S_{n+1}(\omega))^2 d\mu^{\mathbb{N}}(\omega).$$

Furthermore,

$$\begin{aligned} \mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_{n+1}(\omega)F_{f_k}(S_n(\omega)))] &= \mathbb{E}^{\mu^{\mathbb{N}}} \left[\mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_{n+1}(\omega))F_{f_k}(S_n(\omega)) \mid \mathcal{F}_n] \right] \\ &= \mathbb{E}^{\mu^{\mathbb{N}}} \left[F_{f_k}(S_n(\omega)) \mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_{n+1}(\omega)) \mid \mathcal{F}_n] \right] \\ &= \mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_n(\omega))^2], \end{aligned}$$

where we have used the law of total expectation in the first equality. We have shown that

$$I_n = \mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_{n+1}(\omega))^2] - \mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_n(\omega))^2].$$

Therefore, we obtain a telescopic sum

$$\begin{aligned} I &= \lim_{N \rightarrow \infty} \sum_{n=1}^N I_n \\ &= \lim_{N \rightarrow \infty} \mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_{N+1}(\omega))^2] - \mathbb{E}^{\mu^{\mathbb{N}}} [F_{f_k}(S_1(\omega))^2] \\ &= \lim_{N \rightarrow \infty} \mathbb{E}^{\mu^{\mathbb{N}}} \left[\left(\int f_k(S_{N+1}(\omega)x) d\nu(x) \right)^2 \right] - \mathbb{E}^{\mu^{\mathbb{N}}} \left[\left(\int f_k(S_1(\omega)x) d\nu(x) \right)^2 \right] \\ &\leq \|f_k\|_{C^0}^2. \end{aligned}$$

So $I < \infty$ and the series

$$\sum_{n=1}^{\infty} \left(\int f_k(S_n(\omega)Mx) d\nu(x) - \int f_k(S_n(\omega)x) d\nu(x) \right)^2$$

is convergent for $\mu^{\mathbb{N}}$ -a.e. ω and μ -a.e. M . So

$$\begin{aligned} \lim_{n \rightarrow \infty} \int f_k(S_n(\omega)Mx) d\nu(x) &= \lim_{n \rightarrow \infty} \int f_k(S_n(\omega)x) d\nu(x) \\ &= \int f_k(x) d\nu_{\omega}(x). \end{aligned}$$

Since the set ℓ is dense, the result holds for any continuous function, and we have proven the desired result. \square

We now show that the measures ν_{ω} above are necessarily Dirac measures.

Lemma 21. *For $\mu^{\mathbb{N}}$ -a.e. ω , there exists $Z(\omega) \in \mathbb{R}\mathbb{P}^1$ such that $\nu_{\omega} = \delta_{Z(\omega)}$*

Proof. Fix an $\omega \in \Omega^{\mathbb{N}}$ in the full-measure set for which

$$S_n(\omega)_*\nu \Rightarrow \nu_{\omega} \text{ and } S_n(\omega)_*M_*\nu \Rightarrow \nu_{\omega}$$

as $n \rightarrow \infty$ for μ -a.e. M . The sequence $X_n(\omega) = S_n(\omega)\|S_n(\omega)\|^{-1}$ has a convergent subsequence because it is defined on a compact subspace of $\mathbb{R}^{2 \times 2}$. Suppose its limit is $X(\omega)$. As reasoned before,

$$\lim_{n \rightarrow \infty} \frac{S_n(\omega)}{\|S_n(\omega)\|} = X(\omega) \implies \|X(\omega)\| = 1$$

because of the continuity of the norm. Consequently, each $X_n(\omega)$ and $X(\omega)$ itself are non-zero matrices. By Lemma 17, ν is non-atomic, and thus we are in a position to apply Lemma 14 to conclude that

$$X(\omega)_*\nu = X(\omega)_*M_*\nu = \nu_\omega$$

for μ -a.e. M .

Suppose $X(\omega)$ is invertible. This would mean that $\nu = M_*\nu$ and thus X is an element of H_ν as defined in Lemma 15 for μ -a.e. M , therefore $G_\mu \subseteq H_\nu$. We are already assuming that G_μ is closed and have now concluded that it is a subspace of a compact space, which means it must be compact, contradicting assumption (i) of Theorem 9. In conclusion, $X(\omega)$ must not be invertible, from which follows $\text{rank}(X(\omega)) = 1$. By Lemma 8 $X(\omega)_*\nu = \nu_\omega$ is a Dirac measure. \square

4.4. Norm growth. We now prove that convergence to a Dirac measure tells us something about the norm growth of our product of matrices.

Lemma 22. *Let $m \in \mathcal{M}(\mathbb{RP}^1)$ be non-atomic and let (A_n) be a sequence in $\text{SL}_\pm(2, \mathbb{R})$ such that $A_n_*m \Rightarrow \delta_{\bar{z}}$, where $\bar{z} \in \mathbb{RP}^1$. Then $\|A_n\| \rightarrow \infty$. Moreover, for all $v \in \mathbb{R}^2$,*

$$\frac{\|A_n^\top v\|}{\|A_n^\top\|} \rightarrow |\langle v, z \rangle|.$$

Proof. Suppose $A_n\|A_n\|^{-1}$ converges to A . Lemma 14 implies that $\overline{A_n_*m} \Rightarrow \overline{A_*m}$, hence $\overline{A_*m} = \delta_{\bar{z}}$. If $\det A \neq 0$ then $m = \overline{A^{-1}_*\delta_{\bar{z}}}$ is Dirac. Contradiction. Hence $\det A = 0$. Now note that

$$0 = |\det A| = \lim_{n \rightarrow \infty} \left| \frac{\det A_n}{\|A_n\|^2} \right| = \lim_{n \rightarrow \infty} \frac{1}{\|A_n\|^2}$$

so $\lim_{n \rightarrow \infty} \|A_n\| = \infty$ as desired. Furthermore, the fact that $A \neq 0$ tells us that $\text{rank}(A) = 1$ and thus $\text{range}(A)$ is a line. Suppose $\text{range}(A) = \text{span}\{y\} = \bar{y}$ for some $y \in \mathbb{R}^2$, then

$$\overline{A}m(\{\bar{y}\}) = m(\overline{A^{-1}}(\{\bar{y}\})) = m(\mathbb{RP}^1) = 1 = \delta_{\bar{z}}(\{\bar{y}\})$$

and $\bar{z} = \bar{y}$. Now suppose $\|z\| = 1$ and let $\{e_1, e_2\}$ be the canonical basis of \mathbb{R}^2 . Then

$$Ae_1 = \pm\|Ae_1\|z \text{ and } Ae_2 = \pm\|Ae_2\|z.$$

Let

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix},$$

then $\|Ae_1\|^2 + \|Ae_2\|^2 = a^2 + c^2 + b^2 + d^2$. The eigenvalues of $A^\top A$ are $\lambda_1 = 0$ and $\lambda_2 = a^2 + b^2 + c^2 + d^2$ which, together with the fact that $\|A\| = 1$, implies that $\lambda_2 = 1 = \|Ae_1\|^2 + \|Ae_2\|^2$. Now let v be a vector in \mathbb{R}^2 , then

$$\begin{aligned} \|A^\top v\|^2 &= \langle A^\top v, e_1 \rangle^2 + \langle A^\top v, e_2 \rangle^2 \\ &= \langle v, Ae_1 \rangle^2 + \langle v, Ae_2 \rangle^2 \\ &= (\|Ae_1\|^2 + \|Ae_2\|^2) \langle v, z \rangle^2 \\ &= \langle v, z \rangle^2. \end{aligned}$$

Thus,

$$\lim_{n \rightarrow \infty} \frac{\|A_n^\top v\|^2}{\|A_n^\top\|^2} = \langle v, z \rangle^2.$$

□

4.5. Proof of Theorem 9. Define $P_n = M_1^\top \cdots M_n^\top$. Let $\nu \in \mathcal{M}(\mathbb{RP}^1)$ be μ -stationary. By Lemma 19 there exists a measure $\nu_\omega \in \mathcal{M}(\mathbb{RP}^1)$ such that $P_n(\omega)_* \nu \Rightarrow \nu_\omega$ for $\mu^\mathbb{N}$ -a.e. ω . Then, by Lemma 21 there exists a direction $\bar{Z}(\omega) \in \mathbb{RP}^1$ such that $\nu_\omega = \delta_{\bar{Z}(\omega)}$ for $\mu^\mathbb{N}$ -a.e. ω . Using Lemma 22 we obtain that

$$\lim_{n \rightarrow \infty} \|P_n^\top(\omega)\| = \lim_{n \rightarrow \infty} \|P_n(\omega)\| = \infty \quad (4.1)$$

and

$$\frac{\|P_n^\top(\omega)v\|}{\|P_n(\omega)\|} \rightarrow |\langle v, Z(\omega) \rangle|. \quad (4.2)$$

for $\mu^\mathbb{N}$ -a.e. ω and every $v \in \mathbb{R}^2$. Define

$$\begin{aligned} T: \text{SL}_\pm(2, \mathbb{R})^\mathbb{N} \times \mathbb{RP}^1 &\rightarrow \text{SL}_\pm(2, \mathbb{R})^\mathbb{N} \times \mathbb{RP}^1 \\ (\omega, \bar{x}) &\mapsto (\sigma(\omega), M_1(\omega)x) \end{aligned}$$

and

$$\begin{aligned} f: \text{SL}_\pm(2, \mathbb{R})^\mathbb{N} \times \mathbb{RP}^1 &\rightarrow \mathbb{R} \\ (\omega, \bar{x}) &\mapsto \log \frac{\|M_1(\omega)x\|}{\|x\|}. \end{aligned}$$

Then

$$\sum_{j=0}^{n-1} f(T^j(\omega, \bar{x})) = \log \frac{\|M_n(\omega) \cdots M_1(\omega)x\|}{\|x\|} \rightarrow \infty$$

for $\mu^\mathbb{N}$ -a.e. ω and \bar{x} non-orthogonal to $\bar{Z}(\omega)$ by (4.1) and (4.2). Since ν is non-atomic, the convergence holds $\mu^\mathbb{N} \times \nu$ almost everywhere. For

any $w \neq 0$

$$\begin{aligned}
\gamma &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sup_{x \neq 0} \frac{\|P_n^\Gamma x\|}{\|x\|} \right) \\
&\geq \iint \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\frac{\|M_n(\omega) \cdots M_1(\omega) w\|}{\|w\|} \right) d\mu^{\mathbb{N}}(\omega) d\nu(w) \\
&= \int \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} f \circ T^j(\omega, w) d(\mu^{\mathbb{N}} \times \nu)(\omega, w) \\
&= \int f(\omega, w) d(\mu^{\mathbb{N}} \times \nu)(\omega, w)
\end{aligned}$$

by Birkhoff's ergodic theorem. Finally, by Proposition 7

$$\gamma \geq \int f(\omega, w) d(\mu^{\mathbb{N}} \times \nu)(\omega, w) > 0.$$

5. APPLICATION TO CONTINUED FRACTIONS

5.1. Continued fractions and the Lévy constant. It is well-known since Gauss that any irrational number $\alpha \in [0, 1] \setminus \mathbb{Q}$ can be written as a continued fraction in the following way:

$$\alpha = \frac{1}{a_1 + \frac{1}{a_2 + \dots}},$$

where the coefficients $a_1, a_2, \dots \in \mathbb{N}$. This representation is unique, so for each irrational we have an infinite sequence of positive integers a_n . Notice that the coefficients a_n are computed using the Gauss map, $T: [0, 1] \rightarrow [0, 1]$, $T(0) = 0$ and

$$T(x) = \frac{1}{x} \bmod 1, \quad x \neq 0.$$

Indeed,

$$a_n = \lfloor 1/\alpha_{n-1} \rfloor.$$

where we denote the iterates of α under T by

$$\alpha_n := T^n(\alpha), \quad n \in \mathbb{N}_0,$$

and $\lfloor \cdot \rfloor$ represents the integer part of a number. The Gauss map T preserves the absolutely continuous Gauss measure given by

$$g(A) = \frac{1}{\log 2} \int_A \frac{1}{1+x} dx,$$

for any measurable set A , and it is ergodic.

If we truncate the continued fraction expansion up to the coefficient a_n , we obtain a rational number p_n/q_n in irreducible form. The sequence p_n/q_n converges to α with very nice properties (see [12]). These

are called the convergents of α and we have

$$q_n = a_n q_{n-1} + q_{n-2}, \quad p_n = a_n p_{n-1} + p_{n-2}, \quad n \in \mathbb{N},$$

with $q_{-1} = p_0 = 0$ and $q_0 = p_{-1} = 1$. It follows immediately that q_n grows at least exponentially.

The Lévy constant of an irrational α is the exponential growth rate of the inverse of the product of the iterates of α ,

$$\Lambda(\alpha) := \lim_{n \rightarrow +\infty} \frac{1}{n} \log(\alpha_0 \dots \alpha_{n-1})^{-1}.$$

It is not difficult to show that this is greater than zero for any number. It is a remarkable result, shown by Khintchine [14], that this value is constant for a full Lebesgue measure set of numbers. The explicit value was computed by Lévy [15, 16]. We give a simple proof below, based on the fact that we know that T preserves the measure g and using the Birkhoff ergodic theorem. For other measures of interest, in particular, singular measures supported on Cantor sets, this proof would not work.

Theorem 23 (Lévy-Khintchine). *For Lebesgue a.e. $\alpha \in [0, 1] \setminus \mathbb{Q}$,*

$$\Lambda(\alpha) = \frac{\pi^2}{12 \log 2}.$$

Proof. Observe that

$$\Lambda(\alpha) = \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{i=0}^{n-1} \log T^i(\alpha) = \int \log(x) dg(x),$$

for g -a.e. x , where we have used the Birkhoff ergodic theorem. Since g is equivalent to the Lebesgue measure, the above holds also Lebesgue almost everywhere. It remains to compute the integral

$$\int_0^1 \frac{\log(x)}{1+x} dx = - \int_0^1 \frac{\log(1+x)}{x} dx,$$

where we have integrated by parts to obtain the right hand side of the equality. Using the Taylor series of $\log(1+x) = \sum_{k \in \mathbb{N}} (-1)^k x^k / k$ with $|x| < 1$,

$$\Lambda(\alpha) = \frac{1}{\log 2} \sum_{k \in \mathbb{N}} \frac{(-1)^{k+1}}{k^2}.$$

This last series is known to be equal to $\pi^2/12$, as it is a simple exercise using the Fourier series of $x \mapsto x^2$ on $[-\pi, \pi]$ evaluated at 0. \square

The above can be stated in a different context using two dimensional matrices in $G := \mathrm{SL}_{\pm}(2, \mathbb{R})$. Indeed, it is straightforward to check that

$$\begin{bmatrix} \alpha_n \\ 1 \end{bmatrix} = \frac{1}{\alpha_0 \dots \alpha_{n-1}} A_n \dots A_1 \begin{bmatrix} \alpha_0 \\ 1 \end{bmatrix}$$

where

$$A_n = \begin{bmatrix} -a_n & 1 \\ 1 & 0 \end{bmatrix} \in G$$

are hyperbolic. In consequence, we get that the Lévy constant is the Lyapunov exponent

$$\Lambda(\alpha) = \lim_{n \rightarrow +\infty} \frac{1}{n} \log \|A_n \cdots A_1\|.$$

It is also simple to verify that

$$Q_n := A_n \cdots A_1 = (-1)^n \begin{bmatrix} q_n & -p_n \\ -q_{n-1} & p_{n-1} \end{bmatrix}, \quad n \in \mathbb{N},$$

and $Q_0 = I$. In this way, we also have the equality

$$\Lambda(\alpha) = \lim_{n \rightarrow +\infty} \frac{1}{n} \log q_n,$$

giving the exponential growth rate of the convergents of α .

5.2. Random cocycle. Consider the probability measure μ on G defined by

$$\mu \left(\begin{bmatrix} -a & 1 \\ 1 & 0 \end{bmatrix} \right) = p_a$$

for each $a \in \mathbb{N}$, where $0 \leq p_a \leq 1$ and $\sum_{a \in \mathbb{N}} p_a = 1$.

Recall now the shift map $\sigma: G^{\mathbb{N}} \rightarrow G^{\mathbb{N}}$,

$$\sigma(\omega_1, \omega_2, \dots) = (\omega_2, \omega_3, \dots)$$

which is simply the Gauss map T in different coordinates. In addition, take the map $A: G^{\mathbb{N}} \rightarrow G$,

$$A(\omega) = \omega_1.$$

Denote by π the map that transforms a matrix in G into the symmetric of its first entry. Now, take

$$\hat{\pi}(\omega) := (\pi\omega_1, \pi\omega_2, \dots) = (a_1, a_2, \dots).$$

Finally, let

$$\phi(a_1, a_2, \dots) \in [0, 1]$$

be the number whose continued fraction expansion is (a_1, a_2, \dots) (this map ϕ will be further discussed below). Hence, $\phi \circ \hat{\pi}(\text{supp } \mu^{\mathbb{N}}) \subset [0, 1]$.

The linear cocycle (σ, A) over a Bernoulli shift has Lyapunov exponent $\gamma(\omega)$ equal to the Lévy constant $\Lambda(\alpha(\omega))$, where

$$\alpha(\omega) = \phi \circ \hat{\pi}(\omega).$$

Theorem 9 can then be applied.

Proposition 24. $\Lambda(\alpha(\omega)) > 0$ is constant and positive for $\mu^{\mathbb{N}}$ -a.e. ω .

Proof. Easy to check the conditions of Theorem 9. \square

This is a general result depending on the choice of the distribution p_a . In the following we will transform it into a result on irrational numbers.

5.3. Rank intervals. We are now interested in constructing a sort of partition of the interval $(0, 1]$ based on the finite continued fractions of rational numbers. This will allow us in the sequel to relate full probability sets with full Lebesgue measure sets of irrationals.

Given $n \in \mathbb{N}$, define a n -rank interval as a set in the form

$$\Delta_n(k_1, \dots, k_n) = \{\alpha \in [0, 1]: a_1(\alpha) = k_1, \dots, a_n(\alpha) = k_n\}$$

for some integer vector $(k_1, \dots, k_n) \in \mathbb{N}^n$, which is indeed an interval. Moreover, there are only countably many rank intervals. Whenever the integer vector is clear, we may simply write Δ_n . Recall that $a_n(\alpha) = \lfloor 1/\alpha_{n-1} \rfloor$.

We associate the continued fraction expansion of α to the map $\phi: \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{R}$,

$$\phi(a_1, a_2, \dots) := \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\dots}}}$$

As mentioned before, the image of ϕ is $[0, 1] \setminus \mathbb{Q}$ and ϕ is injective. Recall that α is rational iff there is some $n \in \mathbb{N}$ such that $\alpha_n = 0$. So, $a_{n+1}(\alpha) = \infty$. We thus obtain the rationals as limits of irrationals and abuse notation to write

$$\begin{aligned} \phi(a_1, \dots, a_n) &:= \lim_{k \rightarrow +\infty} \phi(a_1, \dots, a_n, k, a_{n+2}, \dots) \\ &= \frac{1}{a_1 + \frac{1}{\dots + \frac{1}{a_n}}} \end{aligned}$$

Observe also that

$$\phi(a_1, \dots, a_n, 1) = \phi(a_1, \dots, a_n + 1).$$

Also, for n even and $k' < k$,

$$\phi(a_1, \dots, a_n, k') < \phi(a_1, \dots, a_n, k),$$

and for n odd and $k' < k$,

$$\phi(a_1, \dots, a_n, k) < \phi(a_1, \dots, a_n, k').$$

Now,

$$\Delta_n(a_1, \dots, a_n) = \begin{cases} [\phi(a_1, \dots, a_n), \phi(a_1, \dots, a_n + 1)), & n \text{ even} \\ (\phi(a_1, \dots, a_n + 1), \phi(a_1, \dots, a_n)], & n \text{ odd.} \end{cases}$$

It is clear that

$$\lim_{n \rightarrow +\infty} |\Delta_n| = 0,$$

where $|\cdot|$ stands for the length of the interval.

Lemma 25. *Let $n \in \mathbb{N}$ and $(a_1, \dots, a_n) \in \mathbb{N}^n$. Then the interiors of the sets*

$$\bigsqcup_{k \in \mathbb{N}} \Delta_{n+1}(a_1, \dots, a_n, k)$$

and $\Delta_n(a_1, \dots, a_n)$ coincide. Moreover, $\sqcup_k \Delta_1(k) = (0, 1]$.

Proof. The intervals $\Delta_{n+1}(a_1, \dots, a_n, k)$ are disjoint by construction. Suppose first that n is odd. So,

$$\begin{aligned} \bigsqcup_{k \in \mathbb{N}} \Delta_{n+1}(a_1, \dots, a_n, k) &= \bigsqcup_{k \in \mathbb{N}} [\phi(a_1, \dots, a_n, k), \phi(a_1, \dots, a_n, k+1)) \\ &= [\phi(a_1, \dots, a_n+1), \phi(a_1, \dots, a_n)), \end{aligned}$$

which has the same interior as $\Delta_n(a_1, \dots, a_n)$. Same idea for n even. \square

The push-forward of the $\mu^{\mathbb{N}}$ measure into $[0, 1]$ is denoted by

$$\eta := \phi_* \hat{\pi}_* \mu^{\mathbb{N}}.$$

From the definitions of Δ_n and η we get immediately that

$$\eta(\Delta_n(a_1, \dots, a_n)) = \prod_{i=1}^n p_{a_i}$$

and

$$\eta(\{x\}) = 0, \quad x \in [0, 1].$$

Additionally,

$$\lim_{n \rightarrow +\infty} \eta(\Delta_n) = 0.$$

Proposition 26. *Any interval $J \subset [0, 1]$ is a disjoint union of rank intervals η -mod0.*

Proof. Suppose $J = (\alpha, \beta)$ have rational edges, i.e. $\alpha, \beta \in \mathbb{Q}$. Let $\phi(a_1, \dots, a_n) = \alpha$ and $\phi(b_1, \dots, b_m) = \beta$ their continued fractions. If $n = m$, it means that both α and β are edges of Δ_n intervals. So, J is the union of $b_n - a_n$ rank intervals (minus α and β themselves).

Without loss of generality, assume now that $n > m$. Hence, since both α and β are edges of n -rank intervals, J is again a union of Δ_n 's.

We now consider the case where $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $\beta \in \mathbb{Q}$. Let $(r_n)_{n \in \mathbb{N}}$ be a strictly decreasing sequence in J of rational numbers such that $r_n \rightarrow \alpha$ and define the sets

$$A_1 = (r_1, \beta), A_2 = (r_2, r_1], \dots$$

The union of these sets is clearly disjoint and equal to J and each A_i can be written as a disjoint union of finitely many rank intervals excluding a set of η -measure zero.

We can apply the same reasoning for the case $\alpha \in \mathbb{Q}$ and $\beta \in \mathbb{R} \setminus \mathbb{Q}$. Finally, if α and β are both irrational, we split the interval by picking a rational r such that $J = (\alpha, r] \cup (r, \beta)$ and treating each subinterval separately. \square

5.4. Lévy constant for subsets of irrationals.

Theorem 27. *If $p_a > 0$ for all $a \in \mathbb{N}$, then η is equivalent to the Lebesgue measure λ on $[0, 1]$.*

Proof. We first prove that $\lambda(A) = 0$ implies $\eta(A) = 0$. Suppose $\lambda(A) = 0$ and let $\varepsilon > 0$. By the familiar characterization of Lebesgue-null sets, there exists a sequence of intervals $U_1, U_2, \dots \subseteq [0, 1]$ such that

$$\sum_{i=1}^{\infty} \lambda(U_i) < \varepsilon \text{ and } A \subseteq \bigcup_{i \in \mathbb{N}} U_i.$$

Without loss of generality, we can assume that U_i fits inside one rank interval (otherwise we can split U_i into subintervals which do). Let $\Delta_{\ell(i)}$ be the highest order interval for some vector $(k_1, \dots, k_{\ell(i)})$ for which U_i is a subset of. Thus,

$$\eta(U_i) \leq \eta(\Delta_{\ell(i)}) = \prod_{j=1}^{\ell(i)} p_{k_j}.$$

This last quantity can be made arbitrarily small by decreasing the chosen value of ε . In particular, for any $\delta > 0$ we can find $\varepsilon > 0$ such that it is smaller than $\delta/2^i$. We have shown that for every $\delta > 0$ there exist intervals $U_1, U_2, \dots \subseteq [0, 1]$ such that

$$\eta(A) \leq \sum_{i=1}^{\infty} \eta(U_i) \leq \sum_{i=1}^{\infty} \frac{\delta}{2^i} = \delta,$$

from which $\eta(A) = 0$ follows.

We now prove the converse. Assume $\eta(A) = 0$. The measure η is regular since it is Borel. Hence, for $\varepsilon > 0$ there exists an open set O such that $A \subseteq O$ and $\eta(O) < \varepsilon$. By Proposition 26, the set O is the countable union of disjoint rank-intervals (η -mod 0) denoted by $\Delta^{(i)}$. Hence,

$$\eta(O) = \sum_{i=1}^{\infty} \eta(\Delta^{(i)}) < \varepsilon.$$

Thus we can cover A with rank-intervals of arbitrarily small η -measure. Because $p_a > 0$ for every $a \in \mathbb{N}$, this implies that we can make $\lambda(\Delta^{(i)})$ as small as desired (if some $p_a = 0$, we would have rank intervals with zero η -measure, so we exclude this situation). That is, we decrease the

η -measure by increasing the order of the rank intervals, which means that their length get also smaller. In particular, we can fix $\delta > 0$ and pick $\varepsilon > 0$ such that $\lambda(\Delta^{(i)}) < \delta/2^i$. From this follows that

$$\lambda(A) < \lambda(O) < \delta,$$

and so $\lambda(A) = 0$. □

The previous results imply that Λ is positive and constant Lebesgue almost everywhere, as in Lévy-Khinchine's theorem. It does not give the actual value of the Lévy constant, as that is the result of the pleasant fact that the invariant Gauss measure is explicitly known. However, the application of Furstenberg's theorem can give some insights about Lévy constants for other sets of numbers, of zero Lebesgue measure. As an example, consider the bounded type irrationals, that is the ones that have a bounded sequence of the continued fraction coefficients: $a_n \leq M$ for some fixed $M > 0$. So, let $p_a = 0$ for $a > M$. From Proposition 24 we also know that the Lévy constant is constant for some full $\mu^{\mathbb{N}}$ measure set. This corresponds to a Cantor set in $[0, 1]$.

The above ideas can be also used for another important function, the Khinchine's constant [14],

$$K_0(\alpha) = \lim_{n \rightarrow +\infty} \frac{1}{n} \log(a_1 \dots a_n).$$

Many questions arise about the properties of the functions Λ and K_0 (see e.g. [8, 22]) with respect to α , which become related to questions about the regularity of γ with respect to μ (cf. [3, 7, 6]). We believe that this connection could be useful to enlarge our understanding of number theory with respect to continued fractions, in particular in higher dimensions.

ACKNOWLEDGMENTS

We would like to thank Jairo Bochi for sharing and explaining to us his notes. JLD was partially supported by the project CEMAPRE - UID/MULTI/00491/2019.

REFERENCES

- [1] M. Artin. *Algebra*. Pearson Education, 2011.
- [2] Jairo Bochi. Furstenberg's theorem on products of i.i.d. 2×2 matrices, 2016.
- [3] Carlos Bocker-Neto and Marcelo Viana. Continuity of Lyapunov exponents for random two-dimensional matrices. *Ergodic Theory Dynam. Systems*, 37(5):1413–1442, 2017.
- [4] Philippe Bougerol et al. *Products of random matrices with applications to Schrödinger operators*, volume 8. Springer Science & Business Media, 2012.
- [5] David Damanik. Schrödinger operators with dynamically defined potentials. *Ergodic Theory Dynam. Systems*, 37(6):1681–1764, 2017.

- [6] João Lopes Dias, Pedro Duarte, José Pedro Gaivão, Silvius Klein, Telmo Peixe, Jaqueline Siqueira, and Maria Joana Torres, editors. *New trends in Lyapunov exponents*. CIM Series in Mathematical Sciences. Springer, 2023. NTLE, Lisbon, Portugal, February 7–11, 2022.
- [7] Pedro Duarte and Silvius Klein. *Lyapunov exponents of linear cocycles*, volume 3 of *Atlantis Studies in Dynamical Systems*. Atlantis Press, Paris, 2016. Continuity via large deviations.
- [8] C. Faivre. The Lévy constant of an irrational number. *Acta Math. Hungar.*, 74(1-2):57–61, 1997.
- [9] Harry Furstenberg. Noncommuting random products. *Transactions of the American Mathematical Society*, 108(3):377–428, 1963.
- [10] Harry Furstenberg and Harry Kesten. Products of random matrices. *The Annals of Mathematical Statistics*, 31(2):457–469, 1960.
- [11] Étienne Ghys. The work of Artur Avila. In *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. 1*, pages 47–54. Kyung Moon Sa, Seoul, 2014.
- [12] G. H. Hardy and E. M. Wright. *An introduction to the theory of numbers*. Oxford University Press, Oxford, sixth edition, 2008. Revised by D. R. Heath-Brown and J. H. Silverman, With a foreword by Andrew Wiles.
- [13] Anatole Katok and Boris Hasselblatt. *Introduction to the modern theory of dynamical systems*. Number 54. Cambridge university press, 1995.
- [14] A. Khintchine. Zur metrischen Kettenbruchtheorie. *Compositio Math.*, 3:276–285, 1936.
- [15] P. Lévy. Sur les lois de probabilité dont dependent les quotients complets et incomplets d’une fraction continue. *Bull. Soc. Math. France*, 57:178–194, 1929.
- [16] P. Lévy. Sur le développement en fraction continue d’un nombre choisi au hasard. *Compositio Mathematica*, 3:286–303, 1936.
- [17] Halsey Royden and Patrick Fitzpatrick. *Real Analysis*. Pearson, 2010.
- [18] J Michael Steele. Kingman’s subadditive ergodic theorem. In *Annales de l’IHP Probabilités et statistiques*, volume 25, pages 93–98, 1989.
- [19] Marcelo Viana. *Lectures on Lyapunov exponents*, volume 145. Cambridge University Press, 2014.
- [20] Marcelo Viana and Kjerfve Oliveira. *Foundations of ergodic theory*. Number 151. Cambridge University Press, 2016.
- [21] Divakar Viswanath. Random Fibonacci sequences and the number 1.13198824... *Math. Comp.*, 69(231):1131–1155, 2000.
- [22] Jun Wu. On the Lévy constants for quadratic irrationals. *Proc. Amer. Math. Soc.*, 134(6):1631–1634, 2006.

ISEG, UNIVERSIDADE DE LISBOA, RUA DO QUELHAS 6, 1200-781 LISBOA, PORTUGAL

Email address: martim@aln.iseg.ulisboa.pt

ISEG AND CEMAPRE-REM, UNIVERSIDADE DE LISBOA, RUA DO QUELHAS 6, 1200-781 LISBOA, PORTUGAL

Email address: jldias@iseg.ulisboa.pt